

No. 3.

Chester. E. Blason.

Notes on The Birds of The Island
of Maui

Stenographic Notes.

7 1 2 3 4 5 6

Birds of Maui.

- Larus barrovianus*.
" *californicus*.
" *delawarensis*.
" *franklinii*.
Sterna fuliginosa.
Puffinus newelli.
Phaethon lepturus.
Fregata aquila.
Anas wyvilliana.
Charadrius albirostris.
Spatula clypeata.
Lafila acuta.
Chen hyperboreus.
Branta nigricans.
Nycticorax nycticorax naevius.
Gallinula sandwicensis.
Fulica alai.
Corymophila falcinaria.
Himantopus knudseni.
Zonotrichia acuminata.
Heteractitis incanous.
Limosa lapponica barrowi.
Calidris acuminata.
Numenius tahitiensis.

Field and Working Notes on The Birds taken
on the Island of Maui

By. Chester E. Blasco.

The following facts are to be observed of
every bird and species taken, if possible
when collecting, etc.

1. The style of bird taken, young or old.

a. Name

(b) measurements etc.

2. Habits of the birds in their native
haunts.

a (actions), b (what they do when
hunting for food), c (whether they haunt
tall or low trees), d (open places or thickly
wooded places), e (and any other action or
habit.)

3. Food. What kind, Where found. Whether
plentiful or scarce. A description of their



food.

4. Actions when eating, etc.

5. Date and locality of observation, elevation,
and on what part of the island they are found.

6.

Digitized by the Internet Archive
in 2017 with funding from
CLIR

B.1. Length 12.70 wing tail
tarsus toe, culmen

Depth of beak. Female, Dec 17, 03
at 1800 foot elevation, in low Ohia
trees about 5 ft. high, at 12:10 P.M.

Found above Kaunapali on SW part
of Maui. Near W. J. Davis' mountain
house.

B.2. Length 11.00 cm. Wing 6.50 cm.

Found on ridge on which W. J. Davis
mountain house is situated, at about 1800
foot elevation, above Kaunapali, Maui,
on Dec. 17, 1903.

B. 3. Length wing. tail
tarsus culmen Depth of beak
thickness width of beak, to

Found as above on Dec. 17. 1903.

B. 4. Length 11.30 wing 6.23 tail

tarsus. culmen to

Depth of beak width of beak,

Female, Found above Kaanapali,

Main, on SW part of island at 1800 ft
elevation, on Dec. 17. 1903. In full song.

Chlorodrepanis wilsoni. Feeding off of

bugs on low Ohia trees. On same

ridge as W. Y. Horner's mountain house
is situated.

B. 5. Length 11.40 wing 6.40 tail
to culmen tarsus

Depth of beak, width of culmen

Found near the head of the Oolawa
Gulch, Maui, Decmbr. 18. 1903; at ²⁵⁰ 750 ft.
elevation (about). Feeding on dactyloctenium
bushes, on the worms that live on the bushes.

Female. *Chlorodrepanis wilsoni*.

Color small, and bird in full plumage, as
to full plumage doubtful. At the lower edge
of the woods in the gulch and shrubs.

B. 6. Length 12. cm wing, 6.30. Found at
1800 foot elevation above Paeano, Maui on
Dec. 21. 1903, on ridge to the eastward of W. G.

Harris' mountain house 2 full song. Male

Chlorodrepanis wilsoni. Testicles enlarged

Also saw a bird, which I am sure
is a *Pseudonotus paultropis*, shot
at it but did not get it. (Oolawa Gulch)



B. 7 - length wing

Found at same place as B. 6.

Pituit

1

Isui

10

Anakahi

21

Eliaio

3

Ahahane

10

Geomys mana

" maculatus

10

Oman

4

Luphe cocconi

1

Ahialoa

1

Ahiapolaau (H. Wilson)

2

Palila

4

Hawaiian Quail

1

Rice Bird

1

English Sparrow

1

Alou

1

Bongun birds

2


Isui

1

Hawks

2

83



Charadrius dominicus fulvus.

Anas interpres.

Dophortyx californica.

Phasianus torquatus.

Turtur chinensis.

Asio accipitrinus sandvicensis.

Acridotheres tristis.

Carpodacus mexicanus obscurus.

Munia risoria

Vestiaria coccinea.

Palmeria dolei.

Himatione sanguinea.

Chlorodrepanis wilsoni.

Oreomysia newtoni.

Scoptes ochracea.

Heterohynchus affinis.

Pseudonestor xanthophrys.

Pittacrostria psittacea.

Stenographic Notes.

3. 1. 2. 3. 4. 5. 6.